

Estimaciones del tamaño efectivo de la población en cepas comerciales y de hatchery de salmón coho a nivel mundial

V. Martínez^{1,*}, P. Dettleff², N. Galarce³, C. Bravo¹, J. Dorner¹, R. Iwamoto⁴ y K. Naish⁵

¹ Favet-Inbiogen, Universidad de Chile, Santiago, Chile

² Facultad de Medicina Veterinaria y Agronomía, Universidad de las Américas, Santiago, Chile

³ Escuela de Medicina Veterinaria, Facultad de Ciencias de la Vida, Universidad Andrés Bello, Santiago, Chile

⁴ Centro de Ciencias Pesqueras del Noroeste, Seattle, EE. UU.

⁵ Facultad de Ciencias Acuáticas y Pesqueras, Universidad de Washington, Seattle, EE. UU.

*jvmartine@uchile.cl

Barbato, M., Orozco-terWengel, P., Tapio, M. & Bruford, M. 2015. SNeP: A tool to estimate trends in recent effective population size trajectories using genome-wide SNP data. *Front. Genet.*, 6: 109.

Carrera, N. 2020. Breve historia de la acuicultura y salmonicultura en el sur de Chile (1856–2000). *Territ. Y Reg.*, 3: 36-49.

Corbin, L., Liu, A., Bishop, S. & Woolliams, J. 2012. Estimation of historical effective population size using linkage disequilibrium with marker data. *J. Anim. Breed Genet.* 129: 257-270.

D'Ambrosio, J., Phocas, F., Haffray, P., Bestin, A., Brard-Fudulea, S., Poncet, C., Quillet, E., Dechamp, N., Fraslin, C., Charles, M. & Dupont-Nivet, M. 2019. Genome-wide estimates of genetic diversity, inbreeding and effective size of experimental and commercial rainbow trout lines undergoing selective breeding. *Genet. Sel. Evol.*, 51: 26.

Dettleff, P., Hernandez, E., Partridge, G., Lafarga-De la Cruz, F. & Martinez, V. 2020. Understanding the population structure and reproductive behavior of hatchery-produced yellowtail kingfish (*Seriola lalandi*). *Aquaculture*, 522: 734948.

Di Prinzipio, C., Penaluna, B., Grech, M., Manzo, L., Miserendino, M. & Casaux, R. 2021. Impact of Chaitén Volcano ashfall on native and exotic fish recovery, recolonization, and abundance. *Sci. Total Environ.*, 752: 141864.

Eldridge, W. & Naish, K. 2007. Long-term effects of translocation and release numbers on fine-scale population structure among coho salmon (*Oncorhynchus kisutch*). *Mol. Ecol.*, 16: 2407-2421.

Falconer, D. & Mackay, T. 1996. *Introduction to Quantitative Genetics*; Longman: Essex, UK.

Frankham, R., Ballou, S., Briscoe, D. & Ballou, J. 2002. *Introduction to Conservation Genetics*; Cambridge University Press: Cambridge, UK.

Hayes, B., Visscher, P., McPartlan, H. & Goddard, M. 2003. Novel multilocus measure of linkage disequilibrium to estimate past effective population size. *Genome Res.* 13: 635-643.

Hershberger, W., Myers, J., Iwamoto, R., McAuley, W. & Saxton, A. 1990. Genetic changes in the growth of coho salmon (*Oncorhynchus kisutch*) in marine net-pens, produced by ten years of selection. *Aquaculture*, 85: 187-197.

Hosoya, S., Kikuchi, K., Nagashima, H., Onodera, J., Sugimoto, K., Satoh, K., Matsuzaki, K., Yasugi, M., Nagano, A., Kumagai, A., Ueda, K. & Kurokawa, T. 2018. Assessment of genetic diversity in Coho salmon (*Oncorhynchus kisutch*) populations with no family records using ddRAD-seq. *BMC Res. Notes*, 11: 548.

Iwamoto, R., Towner, R., Myers, J., Hudson, G. & Munsell, P. 2017. Coho salmon broodstock development: A case study of the Domsea Coho salmon (1977 to 2015). *Bull. Jpn. Fish. Res. Educ. Agency*, 45: 107-112.

Martínez, V., Kause, A., Mäntysaari, E. & Mäki-Tanila, A. 2006. The use of alternative breeding schemes to enhance genetic improvement in rainbow trout (*Oncorhynchus mykiss*): I. One-stage selection. *Aquaculture*, 254: 182-194.

Rozas, M.; Enriquez, R. *Piscirickettsiosis and Piscirickettsia salmonis in fish: A review*. *J. Fish Dis.* 2014, 37, 163–188.

Martínez, V., Naish, K., Bravo, C., Cichero, D., Lien, S., Nome, T., Iwamoto, R. & Moen, T. 2018. Development of a high-density genomic platform for Coho salmon and its use in population assessment in Chile and North America, *Proceedings of the World Congress on Genetics Applied to Livestock Production*. 2: 968.

Martínez, V., Dettleff, P., Galarce, N., Bravo, C., Dorner, J., Iwamoto, R. & Naish, K. 2022. Estimates of Effective Population Size in Commercial and Hatchery Strains of Coho Salmon (*Oncorhynchus kisutch* (Walbaum, 1792)). *Animals*, 12: 647.

Myers, J., Hershberger, W., Saxton, A., & Iwamoto, R. 2002. Estimates of genetic and phenotypic parameters for length and weight of marine net-pen reared coho salmon (*Oncorhynchus kisutch* Walbaum). *Aquac. Res.*, 32: 277-285.

Neely, K., Myers, J. & Hard, J. 2012. Comparison of Early Development between a Domesticated Stock of Coho Salmon and Its Parental Stock. *Trans. Am. Fish. Soc.*, 141: 1504-1509.

Saura, M., Caballero, A., Santiago, E., Fernández, A., Morales-González, E., Fernández, J., Cabaleiro, S., Millán, A., Martínez, P., Palaiokostas, B., Kocour, M., Aslam, M., Houston, R., Prchal, M., Bargelloni, L., Tzokas, K., Haffray, P., Bruant, J-B & Villanueva, B. 2021. Estimates of recent and historical effective population size in turbot, seabream, seabass and carp selective breeding programmes. *Genet. Sel. Evol.*, 53: 85.

Withler, R. & Evelyn, T. 1990. Genetic Variation in Resistance to Bacterial Kidney Disease within and between Two Strains of Coho Salmon from British Columbia. *Trans. Am. Fish. Soc.*, 119: 1003-1009.

Winkler, F., Bartley, D. & Díaz, N. 1999. Genetic differences among year classes in a hatchery population of coho salmon (*Oncorhynchus kisutch* (Walbaum, 1792)) in Chile. *Aquaculture*, 173: 425-433.