

Pan-genoma de aislados chilenos de *Renibacterium salmoninarum*: dilucidando la diversidad en el factor de virulencia p57

M. Saldarriaga-Córdoba^{1,2} y R. Avendaño-Herrera^{2,3,4}

¹ Centro de Investigación en Recursos Naturales y Sustentabilidad, Universidad Bernardo O'Higgins, Santiago, Chile

² Centro Fondap, Interdisciplinary Center for Aquaculture Research (INCAR), Universidad Andrés Bello, Viña del Mar, Chile

³ Laboratorio de Patología de Organismos Acuáticos y Biotecnología Acuícola, Facultad de Ciencias de la Vida, Universidad Andrés Bello, Viña del Mar, Chile

⁴ Centro de Investigaciones Marina Quintay (CIMARQ), Universidad Andrés Bello, Quintay, Chile

*reavendano@yahoo.com – ravendano@unab.cl

- Bayliss, S. C., Verner-Jeffreys, D. W., Ryder, D., Suarez, R., Ramirez, R., Romero, J., Pascoe, B., Sheppard, S. K., Godoy, M., & Feil, E. J. (2018). Genomic epidemiology of the commercially important pathogen *Renibacterium salmoninarum* within the Chilean salmon industry. *Microbial Genomics*, 4: 1-12.
- Brynildsrud, O., Feil, E. J., Bohlin, J., Castillo-Ramirez, S., Colquhoun, D., McCarthy, U., Matejusova, I. M., Rhodes, L. D., Wiens, G. D., & Verner-Jeffreys, D. (2014). Microevolution of *Renibacterium salmoninarum*: Evidence for intercontinental dissemination associated with fish movements. *The International Society for Microbial Ecology Journal*, 8(4), 746–756. doi: 10.1038/ismej.2013.186
- Chien, M. S., T. L. Gilbert, C. Huang, M. L. Landolt, P. J. O'Hara, and J. R. Winton. (1992). Molecular cloning and sequence analysis of the gene coding for the 57-kDa major soluble antigen of the salmonid fish pathogen *Renibacterium salmoninarum*. *FEMS Microbiology Letters* 75, 259-265. doi: 10.1016/0378-1097(92)90414-j
- Choo, K. H., & Ranganathan, S. (2008). Flanking signal and mature peptide residues influence signal peptide cleavage. *BMC Bioinformatics*, 9(Suppl 12), S15. doi: 10.1186/1471-2105-9-S12-S15
- Coady, A. M., Murray, A. L., Elliott, D. G., & Rhodes, L. D. (2006). Both *msa* genes in *Renibacterium salmoninarum* are needed for full virulence in bacterial kidney disease. *Applied and Environmental Microbiology*, 72(4), 2672–2678. doi: 10.1128/AEM.72.4.2672-2678.2006
- Costa, S. S., Guimarães, L. C., Silva, A., Soares, S. C., & Baraúna, R. A. (2020). First steps in the analysis of prokaryotic pan-genomic. *Bioinformatics and Biology Insights*, 14. doi: 10.1177/11779 32220 938064
- Evelyn T.P.T. (1977) An improved growth medium for the kidney bacterium and some notes on using the medium. *Bulletin de l'Office International des Epizooties* 87, 511–513
- Herbert, M., Klein, M., Manting, E. H., Driessen, A. J., & Freudl, R. (1999). Temporal expression of the *Bacillus subtilis* *secA* gene, encoding a central component of the preprotein translocase. *Journal of Bacteriology*, 181(2), 493–500. doi:10.1128/JB.181.2.493-500.1999
- Müller, M., Koch, H. G., Beck, K., & Schafer, U. (2001). Protein traffic in bacteria: Multiple routes from the ribosome to and across the membrane. *Progress in Nucleic Acid Research and Molecular Biology*, 66, 107–157.
- O'Farrell, C.L. and Strom, M.S. (1999). Differential expression of the virulence-associated protein p57 and characterization of its duplicated gene *msa* in virulent and attenuated strains of *Renibacterium salmoninarum*. *Diseases of Aquatic Organisms* 38, 115–123. doi: 10.3354/dao038115
- Ordal, E.J. and Earp, B.J. (1956). Cultivation and transmission of the etiological agent of bacterial kidney disease. *Proceedings of the Society of Experimental Biology and Medicine* 92, 85–88. doi: 10.3181/00379727-92-22392
- Rhodes, L. D., Coady, A. M., & Deinhard, R. K. (2004). Identification of a third *msa* gene in *Renibacterium salmoninarum* and the associated virulence phenotype. *Applied and Environmental Microbiology*, 70(11), 6488–6494. doi: 10.1128/AEM.70.11.6488-6494.2004
- Saldarriaga-Córdoba, M. & Avendaño-Herrera, R. (2022). Comparative pan-genomic analysis of 51 *Renibacterium salmoninarum* indicates heterogeneity in the principal virulence factor, the 57 kDa protein. *Journal of Fish Diseases*. doi: 10.1111/jfd.13653.
- Sanders J.E. & Barros M.J. (1986) Evidence by the fluorescent antibody test for the occurrence of *Renibacterium salmoninarum* among salmonid fish in Chile. *Journal of Wildlife Diseases*, 22, 255–257. doi: 10.7589/0090-3558-22.2.255
- Sanders, J. E., & Fryer, J. L. (1980). *Renibacterium salmoninarum* gen. Nov., sp. nov., the causative agent of bacterial kidney disease in salmonid fishes. *International Journal of Systematic Bacteriology*, 30, 496–502. doi: 10.1099/00207 713-30-2-496
- Smith I.W. (1964) The occurrence and pathology of Dee disease. Department of Agriculture and Fisheries for Scotland. *Freshwater Salmon Fisheries Research* 34, 1–12.
- Tuteja, R. (2005). Type I signal peptidase: An overview. *Archives of Biochemistry and Biophysics*, 441(2), 107–111. doi: 10.1016/j.abb.2005.07.013
- Wiens, G. D. (2011). Bacterial kidney disease (*Renibacterium salmoninarum*). In T. K. Patrick & W. D. Bruno (Eds.), *Fish diseases and disorders*, vol. 3; viral, bacterial and fungal infections (pp. 338–374). CAB International.
- Wiens, G. D., Rockey, D. D., Wu, Z., Chang, J., Levy, R., Crane, S., Chen, D. S., Capri, G. R., Burnett, J. R., Sudheesh, P. S., Schipma, M. J., Burd, H., Bhattacharyya, A., Rhodes, L. D., Kaul, R., & Strom, M.S. (2008). Genome sequence of the fish pathogen *Renibacterium salmoninarum* suggests reductive evolution away from an environmental *Arthrobacter* ancestor. *Journal of Bacteriology*, 190, 6970–6982. doi: 10.1128/JB.00721 -08